

Long-Term INTEGRAL and RXTE Observations of the X-Ray Pulsar LMC X-4

© 2005 S.S.Tsygankov^{1*}, A.A.Lutovinov¹

¹ *Space Research Institute, Russian Academy of Sciences, Profsoyuznaya ul. 84/32, Moscow 117810, Russia*

We analyze the observations of the X-ray pulsar LMC X-4 performed by the INTEGRAL observatory and the All-Sky Monitor (ASM) of the RXTE observatory over a wide energy range. The observed hard X-ray flux from the source is shown to change by more than a factor of 50 (from ~ 70 mCrab in the high state to ~ 1.3 mCrab in the low state) on the time scale of the accretion-disk precession period, whose mean value for 1996–2004 was determined with a high accuracy, $P_{prec} = 30.275 \pm 0.004$ days. In the low state, a flare about 10 h in duration was detected from the source; the flux from the source increased by more than a factor of 4 during this flare. The shape of the pulsar's broadband spectrum is essentially invariable with its intensity; no statistically significant features associated with the possible resonance cyclotron absorption line were found in the spectrum of the source.

© 2005 Pleiades Publishing Inc.

Key words: pulsars, neutron stars, X-ray sources.

* E-mail: st@hea.iki.rssi.ru

INTRODUCTION

The X-ray pulsar LMC X-4 in the Large Magellanic Cloud (LMC) (the distance to the object is $d = 50$ kpc) is a high-mass binary with a pulsation period of ~ 13.5 s (Kelley et al. 1983), in which the compact object is eclipsed by its optical companion, an O8 star of the 14th magnitude with a mass of $20 M_{\odot}$ (Chevalier and Ilovaisky 1977), every ~ 1.408 days (Li et al. 1978; Lang et al. 1981; Levine et al. 2000). Levine et al. (2000) provided the following orbital parameters for the binary: $a_x \sin i = 26.333 \pm 0.019$ light seconds, the eccentricity $e < 0.003(2\sigma)$, and the epoch of the zero orbital phase $T_0 = \text{MJD}51110.86571$.

Lang et al. (1981) pointed to the existence of a superorbital period in the binary, ~ 30.5 days, within which the intensity of the source changes by a factor of about 60. These authors also assumed that this effect is produced by the blockage of direct X-rays by a precessing accretion disk tilted with respect to the orbital plane, much as is the case in the binary Her X-1 (Tananbaum et al. 1972). Based on the measurements of almost the same X-ray flux observed from the source during eclipses in its low and high states, Woo et al. (1995) concluded that the intrinsic luminosity of the pulsar is constant, which also confirms the assumption made by Lang et al. (1981). Having analyzed the RXTE and GINGA observations, Paul and Kitamoto (2002) estimated the rate of decrease in the precession period of the accretion disk to be $\dot{P} \sim -2 \times 10^{-5} \text{ s s}^{-1}$.

The flaring activity of LMC X-4 has been widely discussed in the literature (see, e.g., Epstein et al. 1977; Skinner et al. 1980; Kelley et al. 1983; Levine et al. 1991). A flare with a duration from dozens of seconds to dozens of minutes is detected from the source during its high state, on average, once a day. However, there is also evidence (Woo et al. 1995) for the presence of bursts in the low state.

In different periods, the pulsar LMC X-4 demonstrates both a spin-down and a spin-up, suggesting that its period is close to its equilibrium value. In this case, according to the disk-accretion theory (Ghosh and Lamb 1979), a neutron star with an equilibrium period of ~ 13.5 s and an X-ray luminosity of $\sim 4 \times 10^{38} \text{ erg s}^{-1}$ would have a very high magnetic moment, $\sim 10^{31.5} \text{ G cm}^3$ (Woo et al. 1996).

The spectrum of the source above several keV is described by a power law with a high-energy cutoff. The measured absorption column density toward the source (along with the best-fit parameters) does not depend on the phase of the precession period; its value, $N_H \sim 5.5 \times 10^{20} \text{ atoms cm}^{-2}$, is close to the Galactic column density, suggesting the absence of strong internal absorption in the binary (Naik and Paul 2003). Different authors (Levine et al. 1991; Mihara 1995; Woo et al. 1996; La Barbera et al. 2001) obtained several estimates for the presence of a cyclotron absorption line in the source's spectrum in the energy range 19 to 100 keV, but as yet there are no reliable measurements of its intensity and, accordingly, the surface magnetic field of the neutron star.

In this paper, we present the results of our timing and spectral analyses for the pulsar LMC X-4 performed by using INTEGRAL and RXTE observations over a wide (1–100 keV) energy range. Preliminary results were obtained by Lutovinov et al. (2004).

OBSERVATIONS

The INTEGRAL International observatory (Winkler et al. 2003), which carries four scientific instruments that allow the emission from astrophysical objects to be studied over a wide wavelength range (from optical to hard γ rays), was placed in orbit by a Russian Proton launcher on October 17, 2002 (Eismont et al. 2003).

In this paper, we analyze the data obtained with the ISGRI detector of the IBIS gamma-ray telescope (Ubertini et al. 2003) and the JEM-X X-ray monitor of the INTEGRAL observatory (Lund et al. 2003). The ISGRI detector has an effective area of $\sim 960 \text{ cm}^2$ at 50 keV, is effectively sensitive to photons in the energy range 20 to 200 keV (the energy resolution is $\sim 7\%$ at 100 keV), and allows the image of the sky within a $29^\circ \times 29^\circ$ field of view (the full coding zone is $9^\circ \times 9^\circ$) to be reconstructed with a nominal spatial resolution of ~ 12 arcmin (the angular size of the mask element). See Lebrun et al. (2003) for a more detailed description of the detector. The main elements of the two identical JEM-X modules (JEM-X1 and JEM-X2) are coded-mask xenonmethane gas chambers placed at about 3.2 m from the plane of the detectors. Each module of the telescope has the following technical characteristics: the energy range 3–35 keV, the field of view (the full coding zone) $13.2^\circ (4.8^\circ)$ in diameter, and the detector area 500 cm^2 (Lund et al. 2003).

About 500 individual pointings were made as part of the studies of the LMC region performed by the INTEGRAL observatory from January 2 through January 28, 2003, during which the pulsar LMC X-4 was within the field of view of the INTEGRAL instruments. The total observing time for the source was $\sim 10^6$ s; at all pointings, it was in the full coding zone. We also used data from the All-Sky Monitor (ASM) of the RXTE observatory (http://xte.mit.edu/ASM_lc.html) to analyze the long-period characteristics of the source. These data are daily averaged fluxes from the source in the energy range 1.3–12.2 keV.

We constructed an X-ray image and performed a spectral analysis of the IBIS data using the methods described by Revnivtsev et al. (2004) and Lutovinov et al. (2003). An analysis of a large set of calibration observations for the Crab Nebula revealed a systematic error of $\sim 10\%$ in the estimated absolute flux from the source over a wide energy range; the spectral shape was reconstructed with an accuracy as high as 3–5%. We took this into account in our spectral analysis by adding a systematic error of 5%. All of the errors given in this paper are purely statistical. The standard OSA software of version 4.1, which was provided by the INTEGRAL Science Data Center (<http://isdc.unige.ch>), was used for our timing analysis of the source on pulsation period time scales and for our analysis of the JEM-X data.

Figure 1 shows the IBIS 20–60 keV map of the LMC region. This map was constructed over the entire period of the source’s observations in January 2003. Apart from the pulsar LMC X-4 itself, the black hole LMC X-1 and the single pulsar PSR 054069 are detected in the image at a statistically significant level. No emission is detected from the SN 1987A remnant (Shtykovskiy et al. 2005).

TIMING ANALYSIS

The Precession Period

As was said in the Introduction, there is a period of about 30 days in the binary LMC X-4 that is most likely related to the eclipse of the neutron star’s emitting regions by a precessing tilted accretion disk. To determine the exact precession period, we used the ASM/RXTE data obtained from January 1996 through July 2004. The best precession period of the accretion disk was found by the epoch-folding technique (Leahy et al. 1983) to be 30.275 days. We used the following method to estimate the error of this measurement: for each point on the light curve of the source within a 1σ error in the flux, we randomly chose a point that was used as a new flux from the source. In this way, we modeled another light curve for which, in turn, the best period was determined. We obtained a total of about 100 periods whose spread was considered to be a 1σ confidence interval. Such an analysis yielded the following mean precession period of the accretion disk for 1996–2004: $P_{prec} = 30.275 \pm 0.004$ days.

The hard X-ray (20–50 keV) light curve of the source shown in Fig. 2a has a characteristic shape. About 2 days after the beginning of its observations, the pulsar switched to the on state; in another 5 days, its flux increased by a factor of about 6–8 and reached a peak (~ 70 mCrab). The arrows in the figure indicate the switch-on and switch-off times estimated more than 20 years ago by Lang et al. (1981); the switch-on time was calculated by using the ephemerides derived by these authors; the switch-off time was determined by assuming that the high state of the source lasted $\sim 60\%$ of the entire precession period (Lang et al. 1981). We see that the presumed zero phase (switch-on time) of the superorbital cycle is greatly shifted relative to the current INTEGRAL measurements. Note that the same relationship between the durations of the high and low states is also obtained from the INTEGRAL observations of the source (Fig. 2a).

Unfortunately, the switch-on time of the object under study was not observed by the INTEGRAL observatory directly, and we had to use interpolation to determine the zero phase of the precession period. The switch-on time of the pulsar LMC X-4 estimated in this way is MJD 52644.5 ± 1.0 . Owing to good statistics and a long series of observations, we were able to determine the mean precession period with a high accuracy (see above), which ensures the determination of the presumed zero phase 20 years ago with an accuracy of ~ 1 day by assuming the period to be constant. Using our value of the precession period, we can calculate the presumed zero phase closest to that measured by Lang et al. (1981). Our result is either ~ 18.5 days before or ~ 12 days after this time. Thus, the precession period of the accretion disk, on average, must either increase with $\dot{P} \sim 1.3 \times 10^{-3} \text{ s s}^{-1}$ or decrease with $\dot{P} \sim -2 \times 10^{-3} \text{ s s}^{-1}$. Here, we give lower limits on the rate of change in the period, since we disregard the change that is a multiple of an integer number of periods.

On the other hand, the mean rate of change in the precession period can be estimated by using our values and those obtained by Lang et al. (1981). This estimation yields a value of $\dot{P} = (-2.5 \pm 0.8) \times 10^{-5} \text{ s s}^{-1}$, which is much lower than our previous estimate obtained by extrapolating the switch-on times and agrees, within the error limits, with the

value determined by Paul and Kitamoto (2002).

The observed discrepancy in the estimated rates of change in the precession period may stem from the fact that the precession period of the accretion disk for the pulsar LMC X-4 has evolved not uniformly, but with spinups/spindowns much higher than the average level.

Using long-term ASM/RXTE observations (January 1996–August 2002), Klarkson et al. (2003) constructed a dynamic power spectrum for the source under study. Throughout the period of observations, the precession period was constant near 30.28 days. However, due to the 3σ error of ± 0.46 days, no firm conclusion about the precession-period variations can be reached.

Figure 2b shows the averaged light curve of the pulsar LMC X-4 constructed from ASM/RXTE data over ~ 8.5 years of its observations and folded with the best period of 30.275 days. Interestingly, there is a statistically significant rise in the source’s intensity with a duration of about three days near phase ~ 0.8 of this cycle. A similar feature in the light curve of the pulsar can be seen in Figs. 1 and 3 from Lang et al. (1981). A similar rise in the source’s intensity during the off state is also observed in another X-ray pulsar, Her X-1 (Jones and Forman 1976).

The Orbital Period

Figure 3a shows the 20–50 keV light curve for the pulsar LMC X-4 when the source was near the peak of its observed X-ray flux. The times of the eclipses of the X-ray source by its optical companion are clearly seen from this figure. The ingress and egress were accompanied by smooth variations in the pulsar’s observed intensity. However, a significant rise in the flux from the source with a duration of about three hours was observed a short time (of the order of an hour) after the egress. The ingress and egress duration can be estimated from the source’s light curve, ~ 3 ks. At the neutron star’s orbital velocity of about 500 km s^{-1} (Kelley et al. 1983), this corresponds to the transition region’s size of $\sim 1.5 \times 10^6 \text{ km}$, which is comparable to the sizes of the atmospheres of late-O main-sequence stars (Woo et al. 1996).

The time dependence of the hardness, which is defined as the ratio of the fluxes from the source in the 40–60 and 20–40 keV energy bands, is shown in Fig. 3b. We see a tendency for the radiation from the object under study to soften as it approaches an eclipse, but this dependence breaks down immediately after the eclipse. The hardness during the eclipses themselves is not shown in the figure, because no statistically significant flux can be recorded from the source.

The dashed lines indicate the eclipse ephemerides taken from the papers by Li et al. (1978), Lang et al. (1981), and Levine et al. (2000). Note that the orbital period in the binary is highly stable and has been almost constant over more than twenty years of the pulsar’s studies. The INTEGRAL observations of the source in 2003 confirm this conclusion (see Fig. 3). Several authors (Woo et al. 1996; Levine et al. 2000) have attempted to measure the rate of change in the orbital period, but the significance of the values obtained is low ($\dot{P}_{orb}/P_{orb} = (-5.3 \pm 2.7) \times 10^{-7} \text{ yr}^{-1}$ and $\dot{P}_{orb}/P_{orb} = (1.1 \pm 0.8) \times 10^{-6} \text{ yr}^{-1}$, respectively)

and is consistent with the observational data.

Note that the low-state flux from the source does not drop to zero and is recorded by the INTEGRAL instruments, although it decreases. Thus, for example, according to the observational data spanning the time interval MJD 52662.5–52665, the 20–50 keV flux from the source was 2.6 ± 0.4 mCrab, while several days later, MJD 52667–52668, it decreased by almost a factor of 2, 1.3 ± 0.5 mCrab.

The intensity of the pulsar LMC X-4 decreases sharply during its X-ray eclipses. We analyzed the flux during eclipses for two time intervals, when the source was in its high and low states (MJD 52648.8 and MJD 52665, respectively). In both cases, no emission was detected from the object under study at a statistically significant level, and we obtained only upper limits on its 20–50 keV flux, which proved to be identical, ~ 1.2 mCrab (1σ). Note that during the X-ray eclipse on January 20, 2003, (MJD 52659.94–52660.17), when the source’s intensity decreased, its observed 20–50 keV flux was significantly higher than the above upper limits, 3.7 ± 1.0 mCrab, but the significance of this measurement is low. A similar analysis for the soft (0.1–2.4 keV) X-ray emission was performed by Woo et al. (1995). The results obtained by these authors suggest that the source’s intensity is constant during its eclipses, which they interpreted as evidence that the pulsar’s intrinsic luminosity is constant. This conclusion was based on the assumption that the X rays observed during an eclipse are scattered by the widely distributed matter around the source, possibly the corona above the outer edges of the accretion disk, much as is the case in the pulsar Her X-1 (Lutovinov et al. 2000).

Flares

Many authors have described the flaring activity of the pulsar LMC X-4. Short flares (up to 1000 s in duration), during which the pulsar’s intensity changes by a factor of 2 to 5, are most characteristic of the source under study (Levine et al. 2000).

An rise in the observed flux from the source was recorded during its low state (MJD 52666) on a time scale atypical of it. The flare duration was about 10 h (orbital phase 0.3–0.6), and the peak intensity (~ 10 mCrab) exceeded the average low-state level by a factor of about 4. Figure 4 shows the flare profile constructed from IBIS data in the 20–50 keV energy band.

SPECTRAL ANALYSIS

In the previous section, we showed that a significant number of features could be distinguished in the light curve of the pulsar LMC X-4. Therefore, studying the spectral evolution of the source with time is of considerable interest. Another goal of the spectral analysis is the search for the possible cyclotron resonance absorption line whose existence was mentioned by several authors (see the Introduction).

A power law with an exponential high-energy cut-off (White et al. 1983), which is typical of this class of objects, was chosen as a basis for the spectral analysis of the source over a wide (4–100 keV) energy range:

$$I(E) = A E^{-\alpha} \times \begin{cases} 1, & E < E_c; \\ \exp [-(E - E_c)/E_f], & E \geq E_c, \end{cases} \quad (1)$$

where E is the photon energy in keV, A is the normalization of the power-law component, α is the photon spectral index, E_c is the cutoff energy, and E_f is the e -folding energy in the source’s spectrum. We identified a total of four states in which an independent analysis of the pulsar’s spectrum was performed: the high state (without including the X-ray eclipse times), the bursts immediately after the source’s egress, the low state (without including the X-ray eclipse times), and the flare occurred on January 27, 2003 (MJD 52666).

The best-fit parameters for the source’s spectra based on model (1) are given in the table. We were able to reconstruct the spectra over a wide energy range (using JEM-X data) only in the high state. Since the JEM-X monitor is not sensitive enough, the source was detected at a statistically significant level only by the IBIS telescope. Therefore, when fitting the spectra in this state, we fixed the parameters in the standard X-ray energy range (< 20 keV) at the values obtained for the high state. We see from the table that the shape of the source’s spectrum and its parameters (slope and cutoff parameters) remain almost constant with the pulsar’s intensity variations. Figure 5 shows the energy spectrum of LMC X-4 reconstructed from INTEGRAL data in a wide energy range for the high state.

To test the hypothesis about the presence of a cyclotron feature in the source’s spectrum within the energy range 4–100 keV, we modified the fitting model by adding the corresponding component:

$$\exp \left[-\frac{A_{cyc} W_{cyc}^2 (E/E_{cyc})^2}{(E - E_{cyc})^2 + W_{cyc}^2} \right], \quad (2)$$

where E is the photon energy in keV, E_{cyc} is the cyclotron energy, W_{cyc} is the width of the cyclotron line, and A_{cyc} is its depth.

Using the modified model, we fitted the pulsar’s spectrum during the high state, except the X-ray eclipse times. We used the same procedure that was employed to study the X-ray pulsar KS 1947+300 (Tsygankov and Lutovinov 2005): the energy of the center of the presumed cyclotron line E_{cyc} was varied over the range 5–100 keV at 5-keV step, while its width was fixed at 5 keV; for each trial energy of the line center, we tested the significance of an improvement in statistics using the $\Delta\chi^2$ test. As a result, we found no such energy of the cyclotron line E_{cyc} in the energy range under study whose inclusion in the model would lead to an improvement in the quality of the spectral fit by more than 2σ .

The result obtained may suggest that either the energy of the cyclotron line lies outside our energy range (4–100 keV) or the INTEGRAL instruments are not sensitive enough for the cyclotron line to be detected in the spectrum of the source LMC X-4. The relationship between the spin period of the neutron star and its luminosity, assuming that the period is close to its equilibrium value, argues for the hypothesis of a strong magnetic field ($> 10^{13}$ G) in the binary (Woo et al. 1996).

CONCLUSIONS

We presented the results of our long-term timing and spectral analyses for the X-ray pulsar LMC X-4 performed by using the INTEGRAL observations in January 2003 and the long-term (1996–2004) ASM/RXTE observations. We determined the precession period (30.275 ± 0.004 days) averaged over the last ~ 8.5 years with a high accuracy. Since the source’s switch-on times predicted by different authors do not coincide, we showed that this parameter is not constant and most likely varies nonuniformly. The IBIS/INTEGRAL data revealed small bursts that emerged after the egress of the X-ray source. We detected a flare about 10 h in duration during the low state. On the time scales of the orbital period, its value determined more than 20 years ago is stable and satisfies well our observational data. We performed a spectral analysis for different states distinguishable by the object’s intensity. The spectrum of the source is described by a power law with a high-energy cutoff, which is characteristic of this class of objects. To test the hypothesis about the presence of a cyclotron feature in the source’s spectrum within the energy range 4–100 keV, we properly modified the fitting model. This analysis showed that there is no such feature in the source’s spectrum at a confidence level exceeding 2σ . A comparison of the derived constraints with the results of other authors (Woo et al. 1996) is more likely indicative of a strong magnetic field ($> 10^{13}$ G) on the neutron-star surface than a weak field ($< 5 \times 10^{11}$ G).

ACKNOWLEDGMENTS

We wish to thank E.M. Churazov, who developed the methods and software for analyzing the data from the IBIS telescope of the INTEGRAL observatory. This work was supported by the Russian Foundation for Basic Research (project no. 04-02-17276). We used the data retrieved from the High-Energy Astrophysics Archive at the Goddard Space Flight Center of NASA and the data retrieved from the Archive of the INTEGRAL Science Data Center (Versoix, Switzerland) and the Russian INTEGRAL Science Data Center (Moscow, Russia). This study was performed in part during the visits to the INTEGRAL Science Data Center (Versoix, Switzerland), and we are grateful to its staff for hospitality. A.A. Lutovinov is also grateful to the ESA for financial support of some of these visits.

REFERENCES

1. C. Chevalier and S. Ilovaisky, *Astron. Astrophys.* **59**, L9 (1977).
2. Clarkson W.I., Charles P.A., Coe M.I., et al., *Mon. Not. Roy. Astron. Soc.* **343**, 1213 (2003)
3. N. A. Eismont, A. V. Ditrikh, G. Janin, et al., *Astron. Astrophys.* **411**, L37 (2003).
4. A. Epstein, J. Delvaille, H. Helmken, et al., *Astrophys. J.* **216**, 103 (1977).
5. P. Ghosh and F. Lamb, *Astrophys. J.* 234, 296 (1979).
6. C. Jones and W. Forman, *Astrophys. J.* **209**, L131 (1976).
7. R. L. Kelley, J. G. Jernigan, A. Levine, et al., *Astrophys. J.* **264**, 568 (1983).
8. A. La Barbera, L. Burderi, T. Di Salvo, et al., *Astrophys. J.* **553**, 375 (2001).
9. F. L. Lang, A. M. Levine, M. Bautz, et al., *Astrophys. J.* **246**, L21 (1981).
10. D. A. Leahy, R. F. Elsner, and M. C. Weisskopf, *Astrophys. J.* **272**, 256 (1983).
11. F. Lebrun, J. P. Leray, P. Lavo cat, et al., *Astron. Astrophys.* **411**, L141 (2003)
12. A. Levine, S. Rappaport, A. Putney, et al., *Astrophys. J.* **381**, 101 (1991).
13. A. Levine, S. Rappaport, and G. Zojcheski, *Astrophys. J.* **541**, 194 (2000).
14. F. Li, S. Rappaport, and A. Epstein, *Nature* **271**, 37 (1978).
15. N. Lund, S. Brandt, C. Budtz-Jo ergesen, et al., *Astron. Astrophys.* **411**, L231 (2003)
16. A. A. Lutovinov, S. A. Grebenev, M. N. Pavlinsky, and R. A. Sunyaev, *Astron. Lett.* **26**, 691 (2000).
17. A. A. Lutovinov, S. V. Mol’kov, and M. G. Revnivitsev, *Pis’ma Astron. Zh.* **29**, 803 (2003) [*Astron. Lett.* **29**, 713 (2003)].
18. A. A. Lutovinov, S. S. Tsygankov, M. G. Revnivitsev, et al., in *Proceedings of the V INTEGRAL Workshop*, ESA SP, **552**, 253 (2004).
19. T. Mihara, PhD Thesis (Univ. Tokyo, Tokyo, 1995).
20. S. Naik and B. Paul, *Astron. Astrophys.* **401**, 265 (2003).
21. B. Paul and S. Kitamoto, *J. Astrophys. Astron.* **23**, 33 (2002).
22. M. G. Revnivstev, R. A. Sunyaev, D. A. Varshalovich, et al., *Pis’ma Astron. Zh.* **30**, 430 (2004) [*Astron. Lett.* **30**, 382 (2004)]
23. G. K. Skinner, S. Shulman, and G. Share, *Astrophys. J.* **240**, 619 (1980).
24. H. Tananbaum, H. Gursky, E. Kellog, et al., *Astrophys. J.* **174**, L143 (1972).
25. P. E. Shtykovskiy, A. A. Lutovinov, M. R. Gilfanov, and R. A. Sunyaev, *Pis’ma*

Astron. Zh. **31**, 284 (2005) [Astron. Lett. 31, (2005)].

26. S. S. Tsygankov and A. A. Lutovinov, Pis'ma Astron. Zh. **31**, 99 (2005) [Astron. Lett. 31, 88 (2005)].

27. P. Ubertini, F. Lebrun, G. Di Co cco, et al., Astron. Astrophys. **411**, L131 (2003)

28. N. White, J. Swank, and S. Holt Astrophys. J. **270**, 771 (1983).

29. C. Winkler, T. J.-L. Courvoisier, G. Di Co cco, et al., Astron. Astrophys. **411**, L1 (2003)

30. J. W. Woo, G. W. Clark, and A. M. Levine, Astrophys. J. **449**, 880 (1995).

31. J. W. Woo, G. W. Clark, A. M. Levine, et al.), Astrophys. J. **467**, 811 (1996).

Best-fit parameters for the spectrum of LMC X-4 derived from INTEGRAL data (JEM-X + IBIS)^a

State	Luminosity, $\times 10^{37} \text{ erg s}^{-1}$ ^b	α	E_c , keV	E_f , keV	$\chi^2_N(N)$ ^c
High (MJD 52646-52650)	38.7	0.20 ± 0.15	9.1 ± 0.8	11.0 ± 0.6	0.88(128)
Flares (in high state)	47.9	0.44 ± 0.20	8.9 ± 1.0	11.9 ± 0.8	0.82(126)
Low (MJD 52662.5-52668)	1.4	0.2^d	9.1^d	14.8 ± 3.2	1.19(4)
Flare (MJD 52666)	3.1	0.2^d	9.1^d	17.1 ± 5.3	0.41(6)

^a All errors are given at a 1σ level.

^b The 4–100 keV luminosity at an assumed distance to the source of 50 kpc.

^c The χ^2 value normalized to the number of degrees of freedom N .

^d The parameters were fixed at the values obtained for the high state.




Fig. 1: IBIS/INTEGRAL 20–60 keV image of the LMC region containing the X-ray pulsar LMC X-4. The sources detected at a statistically significant level are labeled; the position of the SN 1987A remnant is indicated by the cross.




Fig. 2: (a) Hard X-ray (20–50 keV) light curve for the pulsar LMC X-4 constructed from IBIS/INTEGRAL data. The arrows indicate the switch-on and switch-off times for the source, as estimated by other authors. (b) The light curve for the source constructed from ASM/RXTE data over eight years of observations and folded with the best period derived from these data. The errors correspond to one standard deviation.




Fig. 3: (a) Orbital light curve for the pulsar LMC X-4 constructed from IBIS data during its high state. The dashed lines indicate the ingress and egress times estimated by Lang et al. (1981). (b) Variations of the source's hardness on the time scale of the orbital period. The errors correspond to one standard deviation.




Fig. 4: Flare detected by the IBIS telescope of the INTEGRAL observatory from the pulsar LMC X-4 on January 27, 2003, during the low state. The solid line indicates a Gaussian fit to the flare profile.




Fig. 5: Energy spectrum of the source LMC X-4 during its high state constructed from JEM-X and IBIS data. The dots indicate the experimental spectrum; the solid line represents its power-law fit with a high-energy cutoff.